

## CT Enterography Patient Instructions

-----  
WEEKS or DAYS before your appointment:  
-----

- **Preparation:** You must follow ALL INSTRUCTIONS in this letter. Your doctor has been notified of your appointment.

Ask your doctor right away about **BLOODWORK** you may need before your appointment, or **MEDICATIONS** you may need to change. **If not complete, you will be rebooked.**

- **Take your weight:** Contact Diagnostic Imaging at the number provided above if your weight is over 400lbs.
- **Arrange childcare:** For safety reasons, children cannot accompany you in procedure rooms and cannot be left unattended.
- **Arrange a support person:** An adult support person is welcome to accompany you to your appointment.
- **Know if you're pregnant:** If you're pregnant, consult with your doctor before proceeding with your appointment.
- **Consult your diabetes care provider:** If you're taking diabetes medications or using Insulin Pump Therapy, consult your diabetes care provider before your appointment. [www.ipumpit.ca](http://www.ipumpit.ca)
- **Remove your continuous glucose monitoring device (CGMD):** If you have a CGMD and it is in the area being imaged, the device may be damaged during your appointment. If the device is imaged, you will need to do finger prick readings, along with your CGMD readings, to ensure it is not damaged.
  - Alberta Health Services will not cover the cost of a replacement CGMD.
  - If you do not know if you need to remove your device or cannot follow these instructions, contact Diagnostic Imaging at the number provided above.

-----  
2 DAYS before your appointment:  
-----

- DO NOT have barium 48 hours before your appointment.
- Continue to take prescribed medication, as directed by your doctor.

-----  
HOURS before your appointment:  
-----

### Adults (18 years and older)

#### 6 Hours Before

- Stop all food. DO NOT chew gum. DO NOT eat candy.
- You may continue drinking **clear fluids**.

#### 3 Hours Before

- Stop drinking **clear fluids**.

### Children (17 years and younger)

#### 3 Hours Before

- Stop all food. DO NOT chew gum. DO NOT eat candy.
- You may continue drinking **clear fluids** until you arrive for your appointment.

## What are clear fluids?

**Allowed:** Fluids that you can see through such as water, apple juice, Gatorade™, Kool-Aid™, Pedialyte™, ginger ale, Sprite™, 7UP™, popsicles, black coffee, or tea

**NOT Allowed:** Milk based fluids, broth or soup, energy drinks, fruit flavoured gelatin, Jello™, orange juice.

## BRING to your appointment:

- Change of clothes (below the waist) in case of an accident.
- Current list of medications (prescription and over the counter).
- Any medication you may need during your appointment.
- Healthcare card.
- Picture identification.
- Eyeglasses, hearing aid(s), and mobility aid/device.
- Patients with diabetes - insulin and glucometer
- Green sleeve that includes your Goals of Care (GOC) Designation Order form if you have one.

## DO NOT bring:

- Leave valuables at home, including jewelry or piercings.
- Limit the number of items you bring to your appointment.

## AT your appointment:

- **Consent is required:** A legal representative (parent or alternate decision maker) must be available (in person or by telephone) to provide consent for all patients under the age of 18, or any patients that do not have the capacity to make health care decisions.
- You will be provided with a hospital gown to change into.
- You will be given contrast; this will make parts of your body more visible in images.
  - You will have an intravenous (IV) line placed in your arm, which will be used to inject contrast.
  - You will be given up to 1.5 litres of oral contrast to drink over 1 hour. Inform your ordering doctor if you think you will not be able to drink this much.
- Your appointment will take up to 2 hours.

## AFTER the appointment:

- Drink lots of fluids after your appointment to help flush the contrast out of your body.
- You may NOT be able to return to your normal activities for several hours after your appointment. The oral contrast may cause you to have frequent bowel movements.